



## **Historic, Archive Document**

Do not assume content reflects current scientific knowledge, policies, or practices.



1  
P698 Sd

S. D.—67. [This leaflet is distributed only with the seeds to which it relates.]

## United States Department of Agriculture,

BUREAU OF PLANT INDUSTRY,

New and Rare Seed Distribution,

WASHINGTON, D. C.

### ALFALFA.

Instructions adapted to Michigan, Wisconsin, and Minnesota.

**OBJECT OF THE DISTRIBUTION.**—The distribution of new and rare seeds has for its object the dissemination of new and rare crops, improved strains of staple crops, and high-grade seed of crops new to sections where the data of the Department indicate such crops to be of considerable promise. Each package contains a sufficient quantity for a preliminary trial, and where it is at all practicable the recipient is urged to use the seed for the production of stocks for future plantings. It is believed that if this practice is followed consistently it will result in a material improvement in the crops of the country. Please make a full report on the inclosed blank regarding the results you obtain with the seed.

#### DESCRIPTION.

Alfalfa (*Medicago sativa*) is a deep-rooted, hardy, perennial forage plant, belonging to the family which includes beans, peas, and clover. It occupies the same place in western agriculture that clover fills in the eastern part of the United States. In chemical composition it resembles the clovers, peas, and allied legumes in having a high protein content, but it slightly outranks any of these that are now under cultivation in percentage of both total and digestible protein. Since it is a perennial, it will last a number of years from one sowing unless crowded out by weeds or otherwise destroyed.

#### SOIL REQUIREMENTS.

A deep, fertile, well-drained, nonacid soil, reasonably free from weeds, is required. The crop can be made to grow on soils of almost any texture, but the proportion of failures is apt to be greater on the deep, porous sands or sands underlain by gravel. It is practically useless to sow alfalfa on thin soils where the bedrock approaches the surface, on land underlain with hardpan, or in locations where the subsoil is so compact that the roots can not penetrate it to considerable depths. It is also equally useless to attempt to grow alfalfa on



land where the fluctuating water table comes near the surface. For the purpose of ascertaining the character of the soil and subsoil and also the depth to the water table, numerous borings should be made with a soil auger. In determining the adaptability of a tract of land to alfalfa, this instrument in most cases will be of greater assistance than a chemical analysis of the soil.

Not only should the land have good underdrainage, but the surface should have sufficient slope to carry off the surplus water readily. In fields that are poorly surface drained sheets of ice may form in the winter. They are fatal to alfalfa. Even on considerable slopes ice may form on the surface, but this is a danger that can not be avoided and is least on the slopes. Rich river or creek bottom lands that have a well-established drainage system are admirably suited to the crop. Overflows may do no harm, provided they are not of long duration and come at a time of the year when growth is practically dormant.

#### PRECEDING CROP.

Alfalfa may follow almost any farm crop if sufficient time is taken to prepare the seed bed and rid the soil of weeds. Cultivated crops, such as potatoes, corn, or beans, are admirably suited for this purpose. Alfalfa may also follow small-grain crops, provided the land has previously been treated in such a way as to destroy the weeds. Sowing on sod land is not recommended unless sufficient time can be given for its preparation to decompose the soil and eradicate the weeds before sowing time. More satisfactory results are secured when such land is devoted to some cultivated crop for a year or two before sowing the alfalfa.

#### PREPARATION OF LAND.

Many of the failures to secure a good stand may be traced directly to the improper condition of the seed bed. The aims to be sought in preparing land for alfalfa are a firm, fine seed bed that is free from weeds. Four to six weeks should intervene between the time of plowing and that of sowing, and meantime the land should be given frequent cultivations. Where it is practicable, fall plowing is recommended. In the spring the land should be disked thoroughly and harrowed at intervals of a week to 10 days until the seed is sown. In some cases it is possible to get a crop of early potatoes off in sufficient time for sowing alfalfa the same season. If the potato ground has been well cultivated it will require little further preparation for alfalfa.

#### LIMING.

The use of lime is not essential on all soils in this region, but it is sometimes difficult to tell whether an application of lime is needed before sowing. If, however, there is any doubt concerning this



point, the land should be limed. It may be applied with a manure spreader, a fertilizer distributor, a lime distributor, or by hand. Any method which spreads the lime uniformly and at low cost is satisfactory. It should be applied at least two or three weeks before sowing, in order that it may become thoroughly incorporated with the soil. At least a ton of burned lime to the acre is generally required and larger applications are often necessary on the heavier soils. If ground limestone or ground oyster shell is used, the quantity should be double that of the burned lime. Experiments have shown very little difference in the final results obtained from the different forms of lime. Burned lime will give quicker results, but the ground limestone and ground oyster shell will finally give the same benefit. The essential element in lime in any form is calcium oxid, and it is recommended that the farmer use whichever form of lime is cheapest, based upon the percentage of this substance present. Where the consumer pays the freight, it should be remembered that he will not only have to pay such charges on practically twice as much of the ground limestone as of the burned lime, but will also be to the additional expense of hauling and spreading 2 tons of the former to 1 of the latter in order to obtain the same results.

#### FERTILIZING.

Well-rotted barnyard manure which is at least comparatively free from weed seeds is the most satisfactory fertilizer. It should be spread on the land before plowing, in order that it may become thoroughly incorporated with the soil. Good results also follow from heavy applications to the preceding crop. Where manure is not available, the judicious use of commercial fertilizers will in many cases increase the chances of success in the sowing of alfalfa. An application of 75 to 100 pounds of potash, 350 to 500 pounds of acid phosphate, and 50 to 75 pounds of nitrate of soda per acre is suggested.

#### INOCULATION.

Nitrogen-fixing bacteria should be provided unless the soil is known to be supplied with these organisms. Inoculation with them may best be accomplished by scattering over the area to be seeded soil from a field upon which the crop has previously been grown successfully. The soil should be broadcasted at the rate of 250 to 500 pounds to the acre and harrowed in immediately. It is suggested that the spreading be done on a cloudy day or in the morning, as the sun's rays on the soil are thought to be destructive to the germs. Soil from the roots of sweet-clover plants also will inoculate alfalfa. In using soil as an agency to inoculate the land, care should be taken to avoid the introduction of noxious weeds and fungous diseases. The practice of sowing a small quantity of alfalfa with the regular sowing

of clover each year for a few years before it is intended to devote the land to alfalfa has in some cases apparently resulted in satisfactory inoculation.

Another method which may be used is that of inoculating the seed with artificial culture, a limited quantity of which can be procured from the United States Department of Agriculture free of charge. Full instructions for use accompany each bottle of culture. The combined use of the soil and the artificial culture is recommended where both can be readily obtained.

#### SOWING.

Alfalfa seed should be sown at the rate of 20 to 25 pounds to the acre. It may be drilled or sown broadcast and covered with a smoothing harrow or weeder. A more even stand is secured if the seed be divided and one-half sown each way of the field. Experience has shown that where the seed is drilled less is required per acre than when it is broadcasted. The use of a nurse crop, while highly desirable, is not generally recommended except in the case of early spring sowings, when from 1 to  $1\frac{1}{2}$  bushels of oats or barley will assist in keeping down the weeds and may cause no serious injury to the alfalfa, provided the nurse crop is cut for hay. In some cases, sowing with early canning peas has also given very good results.

The most successful stands are secured from sowing late in June or as soon thereafter as moisture conditions are favorable. It is seldom safe to sow much later than August 15, as the plants may not become thoroughly established before winter sets in. Successful stands are sometimes secured from early spring sowings, though they are not generally to be recommended.

#### TREATMENT OF THE STAND.

In spring sowings, unless the weeds threaten to choke out the young plants, the alfalfa should not be clipped until it is 12 to 15 inches high and commencing to bloom. The cutter bar should be set high, as the alfalfa is likely to be ruined if it is cut low. If the first cutting is light it may be left on the land as a mulch, but if heavy enough to smother the plants it should be removed. Late summer sowings should not be cut until the following season, and mowing should be discontinued early enough to allow a growth of 8 to 10 inches at the end of the summer to catch and hold the snow and protect the crowns of the plants.

If the alfalfa goes through the winter successfully the first cutting may usually be made in June. It should be made when the plants are beginning to bloom, or, better still, when the basal shoots appear. In an average season three cuttings may be expected. If the plants turn yellow the alfalfa should be cut, no matter at what stage of



development. If the stand becomes very thin or patchy, the field should be plowed and resown. Attempts at patching up poor stands have not generally been successful. If the weeds threaten to destroy the alfalfa a modified form of the spring-tooth harrow may be used with fair results. However, there is much doubt with regard to the value of cultivation for keeping down weeds or otherwise improving the broadcast stands.

A top-dressing of well-rotted, weed-free stable manure, applied in the late fall or early winter, preferably with a manure spreader, will not only furnish plant food but will also serve as a protection during the winter months. Where this is not available a light application of a commercial fertilizer consisting chiefly of acid phosphate with a small quantity of potash will tend to increase the growth and lengthen the life of the field. Under no circumstances should the field be pastured during the first two years, and even an old field should be pastured sparingly, if at all.

#### SOME COMMERCIAL VARIETIES OF ALFALFA.

*Common alfalfa.*—Under this designation is included the greater part of the alfalfa grown in the United States, the seed from the various sources frequently being named after the State in which it was produced. Where alfalfa has been grown under certain conditions for a considerable time there is a tendency through elimination to produce a more or less distinct strain, presumably best adapted to the conditions under which it was developed. Thus, for instance, seed from fields that have resulted from several seed generations in Montana and the Dakotas may produce plants that will prove to be somewhat more hardy than those from seed grown farther south. Such northern-grown seed is preferred to the southern-grown seed for the northern part of the United States.

*Turkestan alfalfa.*—Turkestan alfalfa was introduced into the United States from Turkestan in 1898, and during recent years practically all the seed imported into this country has been from that source. This variety, although quite variable, resembles common alfalfa in general characteristics, but as a rule does not produce quite as heavy yields. Selected strains have proved somewhat superior to the ordinary alfalfa both from the standpoint of hardiness and that of drought resistance. The variety as a whole, however, is generally inferior to the alfalfa commonly grown in this country, especially in the humid sections.

*Grimm alfalfa.*—Grimm alfalfa was introduced into this country in 1857 from Baden, Germany, by Wendelin Grimm, of Carver County, Minn. Careful investigations indicate that it owes its superior hardiness to the fact that it is the result of a natural cross

between the common variety and the yellow-flowered alfalfa (*Medicago falcata*) and that by virtue of its being exposed to numerous severe winters the weaker plants were eliminated, leaving only the more hardy ones to perpetuate the strain. Grimm alfalfa does not differ materially in appearance from the ordinary strain, so that the casual observer has difficulty in distinguishing one from the other. While a large percentage of its flowers are of the same color as those of common alfalfa, there are some that represent many shades of violet, yellow, and other hues. The taproots show a tendency to branch, and the crowns are inclined to be low set and spreading, characteristics which undoubtedly are of great importance in rendering the variety resistant to drought. Grimm alfalfa is one of the hardiest, if not the most hardy, of our commercial strains. It is recommended for sections where the winters are especially severe and where little protection is given by snow. In sections where winterkilling is not an important consideration it is not thought to be materially superior to common alfalfa.

PUBLICATIONS AVAILABLE.

For further details regarding alfalfa, see Farmers' Bulletins 339, "Alfalfa," and 757, "Commercial Varieties of Alfalfa," which will be sent free of charge upon application to the Secretary of Agriculture, Washington, D. C.

Approved:

WM. A. TAYLOR,  
*Chief of Bureau.*

SEPTEMBER 20, 1921.





